


SOLUTIONS:

- (i) $\sin 0 = 0$ (ii) $\sin \frac{\pi}{2} = 1$ (iii) $\sin \pi = 0$ (iv) $\sin \frac{3\pi}{2} = -1$ (v) $\sin 2\pi = 0$
 (vi) $\cos 0 = 1$ (vii) $\cos \frac{\pi}{2} = 0$ (viii) $\cos \pi = -1$ (ix) $\cos \frac{3\pi}{2} = 0$ (x) $\cos 2\pi = 1$
 (xi) $\tan 0 = 0$ (xii)ⁱ $\tan \frac{\pi}{2} = \frac{1}{0}$ (xiii) $\tan \pi = 0$ (xiv)* $\tan \frac{3\pi}{2} = \frac{-1}{0}$ (xv) $\tan 2\pi = 0$



^{i*} Note: for $\tan 90^\circ$ and $\tan 270^\circ$, above, we cannot represent a division-by-zero 'value' on the 'codomain' (y -axis), so we draw 'asymptotes', represented by the vertical dashed lines on the graph. Anyway, $\tan 90^\circ$, $\tan 270^\circ$, $\tan 450^\circ$, etc., do not exist on the 'domain', i.e. they are not 'elements' of the set of the 'domain' (A -axis.) We can say that generally for all " $\tan(90^\circ + m(180^\circ))$ ", where $m \in \mathbb{Z}$. Better to write this in radians i.e. for all $A = \frac{\pi}{2} + m\pi$, where $m \in \mathbb{Z}$.

- SMARTPHONE
- COMPUTER
- TABLET-PC
- PRINTED OUT (A4 PAPER)

visit:

<https://projectmathsnotes.ie/>

AND PURCHASE YOUR COPY ONLINE

Project Maths NotesTM for Leaving Cert

- It's an **investment** in the future of any young person
- Prepares you for the most difficult exam **questions**
- Sorts out **common problems** most students have
- Enables the learner to actually **understand** maths
- Download to your **smart device** - study **on the go**



HIGHER
LEVEL

Copyright Notice

All notes are copyright © M. I. Publishing 1433-41. All rights reserved.

These product samples are for promotional purposes only and may not be edited or parsed.

You must be sufficiently licenced to use notes in private tuition, "grinds" classes, or for teaching.

To order, please visit <https://projectmathsnotes.ie/>